

Decomposition and the Gross-Taylor string

**Queen Mary
October 26, 2023**

Eric Sharpe, Virginia Tech

An overview of T. Pantev, ES, arXiv:2307.08729

The purpose of this talk today is to reconcile two different perspectives on two-dimensional pure Yang-Mills theories:

1) Decomposition

(Hellerman, Henriques, Pantev, ES, Ando '06; ...
..., Nguyen, Tanizaki, Unsal '21, ...)

Two-dimensional pure Yang-Mills = \bigoplus_R (Trivial (invertible) QFTs)

2) Gross-Taylor expansion

(Gross, Taylor '93; Cordes, Moore, Ramgoolam '94, ...)

Two-dimensional pure Yang-Mills = target-space field theory of a string field theory

Executive summary:

Decomposition appears to predict a one-form symmetry in the Gross-Taylor string theory.

Plan of the talk:

1) Review decomposition

Focusing on examples of S_n orbifolds & 2d pure YM

2) Gross-Taylor and two puzzles

Logic of Gross-Taylor:

First rewrite pure YM partition function as a sum of S_n orbifolds, then, interpret those orbifolds as branched covers and then as SFT.

We'll see that the S_n orbifolds interlace with decomposition perfectly, but two puzzles arise in the branched covers/SFT interpretation.

3) Proposed resolution

The branched cover/SFT interpretation will also be compatible if the GT string is required to have a novel symmetry.

A short review of decomposition

In $d > 1$ spacetime dimensions,
if a local quantum field theory has a global $(d - 1)$ -form symmetry,
it is equivalent to a disjoint union of other local QFT's,
known in this context as 'universes.'

We call this **decomposition**.

(2d: Hellerman et al '06, ...;
 $d > 2$: Tanizaki-Unsal '19, Cherman-Jacobson '20, ...)



When this happens, we say the QFT 'decomposes.'

Decomposition has been explored in many examples, as I'll quickly review.

Today: understand decomposition in the Gross-Taylor expansion of 2d pure YM.

More on decomposition...

What does it mean for one local QFT to be a sum of other local QFTs?

(Hellerman et al '06)

1) Existence of projection operators

The theory contains topological local operators Π_i such that

$$\Pi_i \Pi_j = \delta_{i,j} \Pi_j \quad \sum_i \Pi_i = 1 \quad [\Pi_i, \mathcal{O}] = 0$$

Operators Π_i simultaneously diagonalizable; state space = $\mathcal{H} = \bigoplus_i \mathcal{H}_i$

Correlation functions:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_m \rangle = \sum_i \langle \Pi_i \mathcal{O}_1 \cdots \mathcal{O}_m \rangle = \sum_i \langle (\Pi_i \mathcal{O}_1) \cdots (\Pi_i \mathcal{O}_m) \rangle = \sum_i \langle \tilde{\mathcal{O}}_1 \cdots \tilde{\mathcal{O}}_m \rangle_i$$

2) Partition functions decompose

$$Z = \sum_{\text{states}} \exp(-\beta H) = \sum_i \sum \exp(-\beta H_i) = \sum_i Z_i$$

(on a connected spacetime)

Decomposition in 2d gauge theories

(Hellerman et al '06)

Example:

S'pose have G -gauge theory, G semisimple, with finite central $K \subset G$ acting trivially.

Statement of decomposition (in this example):

$$\text{QFT}(G\text{-gauge theory}) = \coprod_{\text{char's } \hat{K}} \text{QFT}(G/K\text{-gauge theory w/ discrete theta angles})$$

Example: pure $SU(2)$ gauge theory = sum $SO(3)_+ + SO(3)_-$ pure gauge theories

where \pm denote discrete theta angles (w_2)

Perturbatively, the $SU(2)$, $SO(3)_\pm$ theories are identical
— differences are all nonperturbative.

Decomposition in 2d gauge theories

(Hellerman et al '06)

Example:

S'pose have G -gauge theory, G semisimple, with finite central $K \subset G$ acting trivially.

As discussed previously, has 1-form symmetry (specifically, BK).

Statement of decomposition (in this example):

$$\text{QFT}(G\text{-gauge theory}) = \coprod_{\text{char's } \hat{K}} \text{QFT}(G/K\text{-gauge theory w/ discrete theta angles})$$

Example: pure $SU(2)$ gauge theory = sum $SO(3)_+$ + $SO(3)_-$ pure gauge theories

where \pm denote discrete theta angles (w_2)

$$SU(2) \text{ instantons (bundles)} \subset SO(3) \text{ instantons (bundles)}$$

The discrete theta angles weight the non- $SU(2)$ $SO(3)$ instantons so as to cancel out of the partition function of the disjoint union.

Summing over the $SO(3)$ theories projects out some instantons, giving the $SU(2)$ theory.

Decomposition in 2d gauge theories

(Hellerman et al '06)

Example:

S'pose have G -gauge theory, G semisimple, with finite central $K \subset G$ acting trivially.

As discussed previously, has 1-form symmetry (specifically, BK).

Statement of decomposition (in this example):

$$\text{QFT}(G\text{-gauge theory}) = \coprod_{\text{char's } \hat{K}} \text{QFT}(G/K\text{-gauge theory w/ discrete theta angles})$$

Formally, the partition function of the disjoint union can be written

$$Z = \underbrace{\sum_{\theta \in \hat{K}} \int [DA] \exp(-S) \exp \left[\theta \int \omega_2(A) \right]}_{\text{Disjoint union}} = \int [DA] \exp(-S) \overbrace{\left(\sum_{\theta \in \hat{K}} \exp \left[\theta \int \omega_2(A) \right] \right)}^{\text{projection operator}}$$

where we have moved the summation inside the integral.

This is an interference effect between universes: **multiverse interference**

$$Z = \sum_{\theta \in \hat{K}} \int [DA] \exp(-S) \exp \left[\theta \int \omega_2(A) \right] = \int [DA] \exp(-S) \left(\sum_{\theta \in \hat{K}} \exp \left[\theta \int \omega_2(A) \right] \right)$$

Disjoint union projection operator

Decomposition in 2d gauge theories

(Hellerman et al '06)

One effect is a projection on nonperturbative sectors:


$$\underbrace{\sum_{\theta \in \hat{K}} \int [DA] \exp(-S) \exp \left[\theta \int \omega_2(A) \right]}_{\text{Disjoint union}} = \int [DA] \exp(-S) \left(\overbrace{\sum_{\theta \in \hat{K}} \exp \left[\theta \int \omega_2(A) \right]}^{\text{projection operator}} \right)$$

Disjoint union of
several QFTs / universes

=

'One' QFT with a restriction on
nonperturbative sectors
= 'multiverse interference'

Schematically,
two theories combine to form a distinct third:



Before going on, let's quickly check these claims for pure $SU(2)$ Yang-Mills in 2d.

The partition function Z , on a Riemann surface of genus g , is

(Migdal, Rusakov)

$$Z(SU(2)) = \sum_R (\dim R)^{2-2g} \exp(-AC_2(R)) \quad \text{Sum over all } SU(2) \text{ reps}$$

$$Z(SO(3)_+) = \sum_R (\dim R)^{2-2g} \exp(-AC_2(R)) \quad \text{Sum over all } SO(3) \text{ reps}$$

(Tachikawa '13)

$$Z(SO(3)_-) = \sum_R (\dim R)^{2-2g} \exp(-AC_2(R)) \quad \begin{array}{l} \text{Sum over all } SU(2) \text{ reps} \\ \text{that are not } SO(3) \text{ reps} \end{array}$$

Result: $Z(SU(2)) = Z(SO(3)_+) + Z(SO(3)_-)$ as expected.

(Later we'll review a more extreme decomposition of 2d pure YM, which we'll compare to GT.)

Since 2005, decomposition has been checked in many examples in many ways. Examples:

- GLSM's: mirrors, quantum cohomology rings (Coulomb branch) (T Pantev, ES '05; Gu et al '18-'20)
- Orbifolds: partition f'ns, massless spectra, elliptic genera (T Pantev, ES '05; Robbins et al '21)
- Open strings, K theory (Hellerman et al hep-th/0606034)
- Susy gauge theories w/ localization (ES 1404.3986)
- Nonsusy pure Yang-Mills ala Migdal (ES '14; Nguyen, Tanizaki, Unsal '21)
- Adjoint QCD₂ (Komargodski et al '20)
- Numerical checks (lattice gauge thy) (Honda et al '21)
- Versions in d-dim'l theories w/ (d-1)-form symmetries (Tanizaki, Unsal, '19; Cherman, Jacobson '20)

This list is incomplete; apologies to those not listed.

Applications include:

- Sigma models with target stacks & gerbes (T Pantev, ES '05)
- Predictions for Gromov-Witten theory (checked by H-H Tseng, Y Jiang, E Andreini, etc starting '08)
- Nonperturbative constructions of geometries in GLSMs (Caldararu et al 0709.3855, Hori '11, ...)
- Elliptic genera (Eager et al '20)
- Anomalies in orbifolds (Robbins et al '21) ..., Romo et al '21)

Today: decomposition in the Gross-Taylor string....

Two examples of decomposition will play an important role in this talk:

- 2d pure Yang-Mills (decomposing to invertibles)
- 2d Dijkgraaf-Witten theory

The role of the first is clear:
we're trying to reconcile decomposition of 2d pure Yang-Mills
with its description ala Gross-Taylor.

Now, part of the Gross-Taylor story is a rewriting of the 2d pure YM partition function as a sum of 2d Dijkgraaf-Witten theories, so its decomposition will also play a role.

We'll discuss each in turn.

Example: 2d pure Yang-Mills (decomposing to invertibles)

Recall from (Migdal '75, Drouffe '78, Lang et al '81, Menotti et al '81, Rusakov '90)
that 2d pure Yang-Mills has been solved exactly.

The partition function $Z(\Sigma)$ on a closed Riemann surface Σ of genus p and area A is

$$Z(\Sigma) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2} C_2(R)\right)$$

where

R is an irrep of the gauge group

$C_2(R)$ is the quadratic Casimir of R

How does it decompose?

Example: 2d pure Yang-Mills (decomposing to invertibles)

Recall from (Migdal '75, Drouffe '78, Lang et al '81, Menotti et al '81, Rusakov '90)
that 2d pure Yang-Mills has been solved exactly.

The partition function $Z(\Sigma)$ on a closed Riemann surface Σ of genus p and area A is

$$Z(\Sigma) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2} C_2(R)\right)$$

Decomposes into theories associated with irreps R :

$$Z(\Sigma) = \sum_R Z_R \quad Z_R = (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2} C_2(R)\right)$$

(It can also decompose along center symmetries,
but the decomposition along irreps will be the focus of the rest of this talk.)

How to interpret those constituent theories?...

Example: 2d pure Yang-Mills (decomposing to invertibles)

2d pure YM is a disjoint sum of trivial ('invertible') field theories,

associated to the irreps R :

(Nguyen, Tanizaki, Unsal '21)

$$Z(\Sigma) = \sum_R Z_R \quad Z_R = (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2} C_2(R)\right)$$

The constituent invertible field theories are ~ classical theories, with 1d Fock space (only vacuum), indexed by counterterms:

$$S = \int_{\Sigma} \sqrt{-g} (aR + b) \quad Z = \exp(a\chi(\Sigma) + b \cdot \text{Area})$$

so the universe associated to irrep R (partition function Z_R)

$$\text{has} \quad a(R) = \ln \dim R, \quad b(R) = -\frac{g_{YM}^2}{2} C_2(R)$$

when interpret as invertible field theory. Next: Dijkgraaf-Witten...

Example: 2d Dijkgraaf-Witten theory

This is a fancy name for an orbifold of a point: $[\text{point}/G]$ for G finite

In cases w/o discrete torsion, operators are twist fields associated to conjugacy classes.

Correlation functions: On a Riemann surface Σ of genus p ,

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \frac{1}{|G|} \sum_{s_1, t_1, \dots, s_p, t_p \in G} \delta \left(\mathcal{O}_1 \cdots \mathcal{O}_n \prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1} \right)$$

where
$$\delta(g) = \begin{cases} 1 & g = 1 \\ 0 & g \neq 1 \end{cases}$$

For example, the partition function is

$$Z = \frac{1}{|G|} \sum_{s_1, t_1, \dots, s_p, t_p \in G} \delta \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1} \right) \quad \text{How does it decompose?}$$

Example: 2d Dijkgraaf-Witten theory

This theory also decomposes into a disjoint sum of trivial ('invertible') field theories, associated to the irreps r .

Projection operators P_r exist:
$$P_r = \frac{\dim r}{|G|} \sum_{g \in G} \chi_r(g^{-1}) g$$

This can also be written as a sum over conjugacy classes, but this form is simpler.

These are projection operators in the sense that $P_r P_s = \delta_{r,s} P_r$, $\sum_r P_r = 1$


Correlation functions in the universe associated to irrep r are

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle_r = \langle \mathcal{O}_1 \cdots \mathcal{O}_n P_r \rangle = \frac{1}{|G|} \sum_{s_1, t_1, \dots, s_p, t_p \in G} \delta \left(\mathcal{O}_1 \cdots \mathcal{O}_n \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1} \right) P_r \right)$$

Note
$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \sum_r \langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle_r$$

Next: Gross-Taylor...

Next, we turn to the Gross-Taylor expansion of 2d pure $SU(N)$ Yang-Mills.



They argued that at large N , this is a target-space SFT of some other 2d string theory, via a series expansion of the partition functions.

Let's review. On a closed Riemann surface Σ_T of genus p and area A ,

$$Z(\Sigma_T) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right)$$

Strictly speaking, to get the right large N asymptotics, we need to write irreps R in terms of coupled representations. For sake of time, and b/c it doesn't significantly affect our result, I'll gloss over that step.

Basic strategy: rewrite the sum over $SU(N)$ irrep data, as a sum over S_n 's and S_n irrep data, where n is the num' boxes in Young tableau for irrep R , and then interpret in terms of branched covers of Σ_T

$$Z(\Sigma_T) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right)$$

Let's rewrite in terms of irreps & characters of the finite symmetric group S_n

Expand the terms using Schur-Weyl duality:

$$\begin{array}{c} \text{\textit{SU}(N) data} \\ \text{(fixed irrep } R) \end{array} \longrightarrow (\dim R(Y))^m = \left(\frac{N^n \dim r(Y)}{|S_n|} \right)^m \frac{\chi_{r(Y)}((\Omega_n)^m)}{\dim r(Y)} \longleftarrow \begin{array}{c} \text{\textit{S}_n} \text{ data} \end{array}$$

where

Y = Young tableau associated with $SU(N)$ irrep R

n = num' boxes in Young tableau Y

$r(Y)$ = S_n irrep associated to Y (and hence $R = R(Y)$)

$$\Omega_n = \sum_{\sigma \in S_n} N^{K_\sigma - n} \sigma$$

K_σ = num' cycles in the cycle decomposition of $\sigma \in S_n$

$$Z(\Sigma_T) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right)$$

$$(\dim R(Y))^m = \left(\frac{N^n \dim r(Y)}{|S_n|}\right)^m \frac{\chi_{r(Y)}((\Omega_n)^m)}{\dim r(Y)}$$

Use the identity $\sum_{s,t \in G} \chi_r(sts^{-1}t^{-1}) = \left(\frac{|G|}{\dim r}\right)^2 \dim r$ to show

$$\begin{aligned} (\dim R(Y))^m &= N^{nm} \left(\frac{\dim r(Y)}{|S_n|}\right)^{m+2p} \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\chi_r\left((\Omega_n)^m \prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right)}{\dim r(Y)} \\ &= N^{nm} \left(\frac{\dim r(Y)}{|S_n|}\right)^{m+2p-1} \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta\left((\Omega_n)^m \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right) P_{r(Y)}\right)}{\dim r(Y)} \end{aligned}$$

One more step....

$$Z(\Sigma_T) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right)$$

So far:

$$(\dim R(Y))^m = N^{nm} \left(\frac{\dim r(Y)}{|S_n|}\right)^{m+2p-1} \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta\left((\Omega_n)^m \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right) P_{r(Y)}\right)}{\dim r(Y)}$$

Use the identity

$$\frac{C_2(R(Y))}{N} = n + \frac{2 \chi_{r(Y)}(T_2)}{N \dim r(Y)} - \frac{n^2}{N^2}$$

to write

$$\begin{aligned} & (\dim R(Y))^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right) \\ &= N^{n(2-2p)} \left(\frac{\dim r(Y)}{|S_n|}\right) \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta\left((\Omega_n)^{2-2p} \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right) P_{r(Y)}\right)}{\dim r(Y)} \exp\left(-g_{YM}^2 \frac{A}{2} n\right) \end{aligned}$$

+ subleading

Finally, we have the Gross-Taylor series expansion.

The partition function of two-dimensional pure $SU(N)$ Yang-Mills

$$Z(\Sigma_T) = \sum_R (\dim R)^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right)$$

has now been rewritten in terms of S_n 's and S_n irrep data:

$$\begin{aligned} & (\dim R(Y))^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right) \xrightarrow{\substack{SU(N) \text{ data} \\ \text{(fixed irrep } R)}} \\ &= N^{n(2-2p)} \left(\frac{\dim r(Y)}{|S_n|}\right) \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta\left((\Omega_n)^{2-2p} \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right) P_{r(Y)}\right)}{\dim r(Y)} \exp\left(-g_{YM}^2 \frac{A}{2} n\right) \\ & \qquad \qquad \qquad \swarrow S_n \text{ data} \\ & \qquad \qquad \qquad + \text{ subleading} \end{aligned}$$

Strictly speaking, we need to break up each irrep R into coupled reps; however, the analysis is nearly identical, and the expression above emerges as one of two chiral components.

Next: interpretation...

Let's interpret:

$$(\dim R(Y))^{2-2p} \exp \left(-g_{YM}^2 \frac{A}{2N} C_2(R) \right) \leftarrow \text{Partition function of a single universe in the decomposition of 2d pure YM.}$$

$$= N^{n(2-2p)} \left(\frac{\dim r(Y)}{|S_n|} \right) \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta \left((\Omega_n)^{2-2p} \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1} \right) P_{r(Y)} \right)}{\dim r(Y)} \exp \left(-g_{YM}^2 \frac{A}{2} n \right) + \text{subleading}$$

The RHS (above) is a sum of 2d Dijkgraaf-Witten correlation functions for group S_n .

In fact, note that the correlation functions have projectors $P_{r(Y)}$
 — these are correlation functions in the universe associated to $r(Y)$!

Takeaway: the partition function of a single universe in the decomposition of 2d pure YM, is a sum of correlation functions in a single universe of 2d Dijkgraaf-Witten for S_n .

Perfect match! Next: Gross-Taylor and 2d strings....

So far: written partition function of a single universe of 2d pure $SU(N)$ Yang-Mills as a sum of correlation functions in a single universe of 2d Dijkgraaf-Witten for S_n

Decomposition meshes perfectly!

Next: interpret in terms of branched covers of the Riemann surface Σ_T

Interpretation of S_n Dijkgraaf-Witten in terms of branched n -covers

(Gross, Taylor '93)

For simplicity, let's take the Riemann surface $\Sigma_T = S^2$


If there are no insertions, then, identify the cover with a disjoint union $\coprod_n S^2$

An insertion of $g \in S_n$ corresponds to a branch point of monodromy g ,
that ties the n sheets of the cover together.

Let's see some examples....

Interpretation of S_n Dijkgraaf-Witten in terms of branched n -covers

Examples: $\Sigma_T = S^2$, $n = 2$: double covers of S^2



S^2 as branched double cover of S^2 ; branch pts at poles, and wraps.

Let's apply to the (original) Gross-Taylor expansion:

$$\begin{aligned} & \sum_R (\dim R(Y))^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right) \\ &= \sum_{n=0}^{\infty} \sum_r N^{n(2-2p)} \left(\frac{\dim r(Y)}{|S_n|}\right) \sum_{s_1, t_1, \dots, s_p, t_p \in S_n} \frac{\delta\left((\Omega_n)^{2-2p} \left(\prod_{i=1}^p s_i t_i s_i^{-1} t_i^{-1}\right)\right)}{\dim r(Y)} \exp\left(-g_{YM}^2 \frac{A}{2} n\right) \\ & \hspace{20em} + \text{subleading} \end{aligned}$$

This is the expansion of the full YM theory — includes sum over all representations (so the projectors $P_{r(Y)}$ sum out — we'll return to them when we look at individual universes).

$$\Omega_n = \sum_{\sigma \in S_n} N^{K_\sigma - n} \sigma$$

Powers of N :

$$\begin{aligned} n(2-2p) + \sum_j \left(K_{\sigma_j} - n\right) &= n\chi(\Sigma_T) + \sum_j \left(K_{\sigma_j} - n\right) \\ &= \chi(\Sigma_W) \quad (\text{Riemann-Hurwitz theorem}) \end{aligned}$$

where Σ_W is a branched n -fold cover of Σ_T

Let's apply to the (original) Gross-Taylor expansion:

$$\begin{aligned} \sum_R (\dim R(Y))^{2-2p} \exp\left(-g_{YM}^2 \frac{A}{2N} C_2(R)\right) \\ = \sum_{n=0}^{\infty} \sum_{s_i, t_i \in S_n} \sum_{L=0}^{\infty} \sum_{v_1, \dots, v_L \in S_n} N^{\chi(\Sigma_W)} (\#) \delta\left(v_1 \cdots v_L \left(\prod_{i=1}^p [s_i, t_i]\right)\right) \exp\left(-\frac{A}{\alpha'_{GT}} n\right) \\ + \text{subleading} \end{aligned}$$

where

$\Sigma_W =$ branched n -fold cover of Σ_T , branched over L points

$$\alpha'_{GT} = \frac{2}{g_{YM}^2}$$

$\# =$ misc' numerical factors, which match Euler char' of space of maps

This is the form expected if 2d pure YM is the SFT of a sigma model $\Sigma_W \rightarrow \Sigma_T$, at large N

Now let's turn to the decomposition.

The partition function of a single universe of 2d pure YM is

$$\begin{aligned}
 & (\dim R(Y))^{2-2p} \exp \left(-g_{YM}^2 \frac{A}{2N} C_2(R) \right) \\
 &= \sum_{s_i, t_i \in S_n} \sum_{L=0}^{\infty} \sum_{v_1, \dots, v_L \in S_n} N^{\chi(\widetilde{\Sigma}_w)} (\#) \delta \left(v_1 \cdots v_L \left(\prod_{i=1}^p [s_i, t_i] \right) \underline{P_{r(Y)}} \right) \exp \left(-\frac{A}{\alpha'_{GT}} n \right) \\
 & \qquad \qquad \qquad + \text{subleading}
 \end{aligned}$$

- Restrict to single $SU(N)$ irrep $R(Y)$
- which fixes $n = \text{num' boxes in Young diagram } Y$ for irrep $R(Y) = \text{covering map deg'}$
- plus added factor of projector $P_{r(Y)}$ in the delta function

This means:

- 1) Sigma model is restricted to maps of a single degree (n)
- 2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

So, we have puzzles to explain in the expansion of a single YM universe:

- 1) Sigma model is restricted to maps of a single degree (n)
- 2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

In broad brushstrokes, both phenomena are typical in decomposition:

- Restrictions on instantons / nonperturbative sectors
- Individual universes can receive contributions which cancel out in sums over universes as we saw previously in the $SU(2) = SO(3)_+ \amalg SO(3)_-$ example.

However, the details here are more extreme:

- Restrictions are usually to a subset of instantons, not to a single instanton degree
- Here the extra contributions would expand possible worldsheets beyond smooth Riemann surfaces

Let's examine in detail...

1) Sigma model is restricted to maps of a single degree (n)

In a 2d NLSM, this is a restriction to (worldsheet) instantons of a single degree.

In decomposition, one often sees restrictions on instanton degrees.

For example, in the $SU(2) = SO(3)_+ \amalg SO(3)_-$ example,
 $SU(2)$ instantons are a subset of $SO(3)$ instantons.


However, in that case, and most other examples,
one restricts to a subset of instantons,
not to instantons of a single degree.

Let's take a moment to review some underlying physics....

Suppose we try to require that the total instanton number always vanish in our QFT.

Start with a field configuration with no net instantons.

Now, move them far away from one another:



Nonzero
instanton number
here!

Total instanton number : 0

Nonzero
instanton number
here!

If physics is local (“cluster decomposition”),
then in those widely-separated regions, the theories have instantons.

So, even if we start with no net instantons,
cluster decomposition implies we get instantons!

Cluster decomposition:



For this reason, Steven Weinberg taught us:

All local quantum field theories must sum over all instantons,
so as to preserve cluster decomposition.

Loophole:

Disjoint unions of QFTs also violate cluster decomposition
(ex: multiple dimension zero operators),
but in principle are straightforward to deal with.

So, if a theory with a restriction on instantons is also a disjoint union,
of theories which are well-behaved, then all is OK.



1) Sigma model is restricted to maps of a single degree (n)

In a 2d NLSM, this is a restriction to (worldsheet) instantons of a single degree.

This is more extreme than we ordinarily see in decomposition.

Furthermore,

labelling field configurations by instanton number
is typically just an artifact of a semiclassical expansion,
and ordinarily does not have an intrinsic meaning in QFT.

Proposal:

the Gross-Taylor string has a symmetry for which map degree is a conserved quantity.

But map degree is a 2-form ($\phi^*\omega$),
so such a symmetry would be either a 1-form or (-1)-form symmetry.

1) Sigma model is restricted to maps of a single degree (n)

Proposal:

the Gross-Taylor string has a symmetry for which map degree is a conserved quantity.

But map degree is a 2-form ($\phi^*\omega$),
so such a symmetry would be either a 1-form or (-1)-form symmetry.

To make this more concrete,
next I'll walk through a related example, where precisely this happens:
2d pure Maxwell theory.

1) Sigma model is restricted to maps of a single degree (n)

2d pure Maxwell theory:

Pure Maxwell theory in any dimension has a global $BU(1)$ (1-form) symmetry:

$$A \mapsto A + \Lambda$$

and Noether current $J^e = *F$, associated to operator $U_\alpha(p) = \exp(i\alpha *F(p))$

In 2d, it also has a magnetic (-1)-form symmetry,

with current $J^m = F$, associated to operator $U_\beta(\Sigma) = \exp\left(i\beta \int_\Sigma F\right)$

So, the symmetries are of the same form as proposed for Gross-Taylor, making it a useful prototype....

1) Sigma model is restricted to maps of a single degree (n)

2d pure Maxwell theory:

$$Z(\Sigma) = \int [DA] \exp(-S) \quad \text{for} \quad S = \frac{1}{g_{YM}^2} \int_{\Sigma} F^{\mu\nu} F_{\mu\nu} + i\theta \int_{\Sigma} F$$

$$\propto \sum_{n=-\infty}^{\infty} \exp\left(-\frac{n^2}{g_{YM}^2 A} + i\theta n\right) \quad \text{where} \quad n \sim c_1 \sim \int F$$

After Poisson resummation,

$$Z(\Sigma) = \sum_{m=-\infty}^{\infty} \exp\left(-\frac{g_{YM}^2 A}{4} (\theta + 2\pi m)^2\right)$$

This is the form of the exact expression for pure YM.

(Paniak, Szabo '02; Gross, Matytsin, '94;
Minahan, Polychronakos, '93;
Caselle et al '93; Fine '90)

Decomposes into universes indexed by m (irreps of $U(1)$), *Poisson dual* to $n \sim c_1$.

1) Sigma model is restricted to maps of a single degree (n)

2d pure Maxwell theory:

$$Z(\Sigma) = \sum_{m=-\infty}^{\infty} \exp\left(-\frac{g_{YM}^2 A}{4} (\theta + 2\pi m)^2\right) \propto \sum_{n=-\infty}^{\infty} \exp\left(-\frac{n^2}{g_{YM}^2 A} + i\theta n\right)$$

Decomposes into universes indexed by m (irreps of $U(1)$), *Poisson dual* to $n \sim c_1$.

Partition function of a single universe is $\exp\left(-\frac{g_{YM}^2 A}{4} (\theta + 2\pi m)^2\right)$

Analogue of the Witten effect:

Shifting $\theta \mapsto \theta + 2\pi$ is equivalent to changing the universe: $m \mapsto m + 1$

1) Sigma model is restricted to maps of a single degree (n)

2d pure Maxwell theory:

$$Z(\Sigma) = \sum_{m=-\infty}^{\infty} \exp\left(-\frac{g_{YM}^2 A}{4} (\theta + 2\pi m)^2\right) \propto \sum_{n=-\infty}^{\infty} \exp\left(-\frac{n^2}{g_{YM}^2 A} + i\theta n\right)$$

This is a prototype for the Gross-Taylor proposal:
there's a decomposition, into universes indexed by m ,
which is Poisson dual to the bundle degree.

In Gross-Taylor, we propose there exists a symmetry which allows us to pick out sectors of
single map degree (single worldsheet instanton number),
which is analogous.

1) Sigma model is restricted to maps of a single degree (n)

So far, we've proposed that the Gross-Taylor string admits an extra symmetry.

Can that be seen directly?

There are (at least) 2 proposals in the literature for the Gross-Taylor string:

1) Cordes-Moore-Ramgoolam: GT string = modification of A model TFT

Standard kinetic terms; localizes on holomorphic maps $\{\bar{\partial}x = 0\}$

2) Horava: GT string = twisted NLSM with nonstandard kinetic terms

Localizes on harmonic maps $\{\partial\bar{\partial}x = 0\}$

The desired symmetry is not immediately visible in either;
might be realized nonlinearly, or, maybe there exists a third version.

Review: puzzles to explain in the expansion of a single YM universe:

1) Sigma model is restricted to maps of a single degree (n)

We've argued this implies the GT string has a new symmetry.

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

We'll study this problem next.

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

Example: $\Sigma_T = S^2$ ($p = 0$), $n = 2$

$$\begin{aligned}
 Z &= \frac{N^{2n}}{n!} \delta \left((\Omega_n)^2 P_r \right) = \frac{N^{2n}}{n!} \delta \left((1)P_r + 2 \left(\frac{1}{N} \right) vP_r + \left(\frac{1}{N} \right)^2 v^2 P_r \right) \\
 &= \frac{N^4}{2!} \delta (P_r) + 2 \frac{N^3}{2!} \delta (vP_r) + \frac{N^2}{2!} \delta (v^2 P_r) \\
 &= \frac{N^4}{4} \pm \frac{N^3}{2} + \frac{N^2}{4}
 \end{aligned}$$

$\Sigma_W = S^2 \amalg S^2$	$????$	$\Sigma_W = S^2$
$\chi(\Sigma_W) = 4$		$\chi(\Sigma_W) = 2$

The N^3 term is new — not present in original GT — present here only b/c of P_r .

How to interpret? $N^\chi = N^3$ so $\chi = 3$, but no closed string worldsheet has χ odd

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

How to interpret? No closed string worldsheet has χ odd

Some options:

- Expand out the projector P_r

In the previous example, we'd get a term prop' to $N^3 \delta(vv)$.
From the delta, should be S^2 , but wrong Euler characteristic.

- Open string?

Subleading corrections were interpreted in the old literature as nonpert' corrections;
open string worldsheets could have odd χ

But these terms aren't all subleading, so expect them to be perturbative,
hence not from open worldsheets.

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

How to interpret? No closed string worldsheet has χ odd

Another possible option: stacky worldsheets

Returning to previous example ($\Sigma_T = S^2, n = 2$):

$$\begin{aligned} Z &= \frac{N^4}{2!} \delta(P_r) + \underline{2 \frac{N^3}{2!} \delta(vP_r)} + \frac{N^2}{2!} \delta(v^2P_r) \\ &= \frac{N^4}{4} \pm \underline{\frac{N^3}{2}} + \frac{N^2}{4} \end{aligned}$$

Interpret as 2 copies of S^2 with a single \mathbb{Z}_2 orbifold point ($\mathbb{P}_{[1,2]}^1$)

$$\chi\left(\mathbb{P}_{[1,2]}^1\right) = 3/2 \quad \chi\left(\mathbb{P}_{[1,2]}^1 \amalg \mathbb{P}_{[1,2]}^1\right) = (2)(3/2) = 3$$

matches power of N !

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

How to interpret? No closed string worldsheet has χ odd

Another possible option: stacky worldsheets

For $\Sigma_T = S^2$, there is a systematic construction of stacky Σ_W 's
(here, Riemann surfaces w/ orbifold points)
that gives matching powers of N .

Idea: Given $\delta(v_1 \cdots v_L)$, write each $v_i \in S_n$ as a product of cycles.
On j th copy of S^2 , if j appears in a cycle of length k , insert \mathbb{Z}_k

Example: S'pose $n = 6$ and $v = (12)(345)(6)$

Then, insert \mathbb{Z}_2 on 2 copies, \mathbb{Z}_3 on 3 copies, smooth pt on last copy.

Can show $\chi = n(2 - 2p) + \sum_j (K_{v_j} - n)$ which matches power of N

2) Presence of projector $P_{r(Y)}$ implies add'l contributions not present previously

How to interpret? No closed string worldsheet has χ odd

Another possible option: stacky worldsheets

Issues:

- Construction only understood for S^2 , not higher genus
- Construction not unique — orb' points can be redistributed across sheets of cover
- Have not tried to compare Hurwitz moduli spaces in general cases

In the same spirit, at least on $\Sigma_T = S^2$,
one can reinterpret the terms as contributions from `stacky' copies of Σ_T ,
meaning, copies with orbifold points.

This is in the spirit of the decomposition:
instead of a sigma model summing over maps $\Sigma_W \rightarrow \Sigma_T$,
this would reflect a decomposition, to trivial field theories
(corresponding to copies of Σ_T).

Summary: reconciling decomposition & GT string pictures of 2d pure YM

1) Reviewed decomposition

Focusing on examples of S_n orbifolds & 2d pure YM

2) Gross-Taylor and the puzzles

Logic of Gross-Taylor:

First rewrote pure YM partition function as a sum of S_n orbifolds, then, interpreted those orbifolds as branched covers and then as SFT.

We saw that the S_n orbifolds interlace with decomposition perfectly, but two puzzles arise in the branched covers/SFT interpretation.

3) Proposed resolution

The branched cover/SFT interpretation will also be compatible if the GT string is required to have a novel symmetry.

Thank you for your time!