

Chiral rings in $2d$ $(0,2)$ theories

Eric Sharpe
Virginia Tech

B Jia, ES, J Guo, to appear

B Jia, ES, R Wu, 1401.1511

R Donagi, J Guffin, S Katz, ES, 1110.3751, 1110.3752

Over the last half dozen years, there's been a *tremendous* amount of progress in perturbative string compactifications.

A few of my favorite examples:

- nonpert' realizations of geometry (Pfaffians, double covers)
(Hori-Tong '06, Caldararu et al '07,...)
- perturbative GLSM's for Pfaffians (Hori '11, Jockers et al '12,...)
- non-birational GLSM phases - physical realization of homological projective duality
(Hori-Tong '06, Caldararu et al '07, Ballard et al '12; Kuznetsov '05-'06,...)
- examples of closed strings on noncommutative res'ns
(Caldararu et al '07, Addington et al '12, ES '13)
- localization techniques: new GW & elliptic genus computations, role of Gamma classes, ...
(Benini-Cremonesi '12, Doroud et al '12; Jockers et al '12, Halverson et al '13, Hori-Romo '13, Benini et al '13,)
- heterotic strings: nonpert' corrections, 2d dualities, non-Kahler moduli (many)

Far too much to cover in one talk! I'll focus on just one....

Today I'll restrict to

- heterotic strings: nonpert' corrections, 2d dualities, non-Kahler moduli

My goal today is to give a survey of some of the results in heterotic strings & $(0,2)$ supersymmetric worldsheets over the last few years, through the lens of chiral rings.

Unfortunately, some topics I won't have the time to discuss: most prominently, non-Kahler moduli.

So, what will I discuss?...

Outline:

- Chiral states in 2d (0,2) NLSM's
- Product structures in chiral rings in $A/2$, $B/2$ twists:
quantum sheaf cohomology
- Nonabelian GLSMs:
 - Dualities in 2d and their geometry
 - Tests of Gadde-Gukov-Putrov triality



Review: chiral rings in 2d (2,2) NLSM's

Consists of states annihilated by
1 of left-moving & 1 of right-moving supercharges.

4 distinct possibilities, labelled (c,c) , (a,c) , (c,a) , (a,a)

In a NLSM on a complex Kahler manifold X ,
all correspond to cohomology of X .

More explicitly...

Review: chiral rings in 2d (2,2) NLSM's

In a (R,R) sector, in a NLSM on a space X , states have the schematic form

$$b_{\bar{i}_1 \cdots \bar{i}_q}^{j_1 \cdots j_p}(\phi) \psi_+^{\bar{i}_1} \cdots \psi_+^{\bar{i}_q} \psi_{-,j_1} \cdots \psi_{-,j_p} |0\rangle$$

ψ_{\pm} worldsheet fermions, $\sim TX$

$$Q = Q_+ + Q_- \leftrightarrow d$$

Q-cohomology classes, counted by $H^{p,q}(X)$

Sit in a topologically protected subsector.

For a (0,2) NLSM, on space X with bundle \mathcal{E} , we'll again look at (R,R) sector states....

For 2d (0,2) NLSM's on Calabi-Yau's (CY's),
Distler-Greene ('88) worked out the analogue:

In a (R,R) sector, zero-energy Q_+ -closed states of form

$$b_{\bar{i}_1 \cdots \bar{i}_q}^{a_1 \cdots a_p}(\phi) \psi_+^{\bar{i}_1} \cdots \psi_+^{\bar{i}_q} \lambda_{-,a_1} \cdots \lambda_{-,a_p} |0\rangle$$

close to large radius.

ψ_+, λ_- worldsheet fermions, $\sim TX, \mathcal{E}$

$$Q_+ \leftrightarrow \bar{\partial}$$

$$\begin{aligned} \text{States counted by } Q_+\text{-cohomology} &= H^q(X, \wedge^p \mathcal{E}^*) \\ &= H^{p,q}(X) \text{ when } \mathcal{E} \cong TX \text{ ((2,2) locus)} \end{aligned}$$

Assumed $K_X, \det \mathcal{E}$ trivial

Q_+ -cohomology no longer in a topological subsector,
but should be protected from perturbative corrections.

So, for large-radius CY, should be reliable.

Consider a more general 2d (0,2) NLSM near large-radius:

$K_X, \det \mathcal{E}$ need not be trivial

The zero-energy Q_+ -closed states again of the form

$$b_{\bar{i}_1 \dots \bar{i}_q}^{a_1 \dots a_p}(\phi) \psi_{+}^{\bar{i}_1} \dots \psi_{+}^{\bar{i}_q} \lambda_{-,a_1} \dots \lambda_{-,a_p} |0\rangle$$

but now $|0\rangle \sim (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2}$

for the Fock vacuum $\psi_{+}^i |0\rangle = 0 = \lambda_{-, \bar{a}} |0\rangle$

States counted by

$$H^q \left(X, (\wedge^p \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right)$$

Different Fock vacua choices give equivalent results....

If instead we'd worked with a Fock vacuum defined by

$$\psi_+^i |0\rangle' = 0 = \lambda_{-,a} |0\rangle'$$

then this one related to last one by

$$|0\rangle' = \left(\prod_a \lambda_{-, \bar{a}} \right) |0\rangle \quad \begin{array}{l} |0\rangle \sim (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \\ |0\rangle' \sim (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \end{array}$$

and states of the form

$$b_{\bar{i}_1 \dots \bar{i}_q}^{\bar{a}_1 \dots \bar{a}_p} (\phi) \psi_+^{\bar{i}_1} \dots \psi_+^{\bar{i}_q} \lambda_{-, \bar{a}_1} \dots \lambda_{-, \bar{a}_p} |0\rangle'$$

$$\begin{aligned} \text{Counted by } & H^q \left(X, (\wedge^p \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \right) \\ &= H^q \left(X, (\wedge^{r-p} \mathcal{E}^*) \otimes (\det \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \right) \\ &= H^q \left(X, (\wedge^{r-p} \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right) \\ & \hspace{15em} \text{(matching previous counting)} \end{aligned}$$

States:

$$H^\bullet \left(X, (\wedge^\bullet \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \right) = H^\bullet \left(X, (\wedge^\bullet \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right)$$

Special case: (2,2) locus

$$\mathcal{E} = TX$$

$$H^\bullet \left(X, (\wedge^\bullet \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right) = H^\bullet (X, \Omega_X^\bullet) = H^{\bullet, \bullet}(X)$$

as expected

On a Calabi-Yau, or if $K_X^{\otimes 2} \cong \mathcal{O}_X$

$$H^\bullet \left(X, (\wedge^\bullet \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \right) = H^\bullet (X, \wedge^\bullet TX)$$

Other tests:

- Invariance under $\mathcal{E} \leftrightarrow \mathcal{E}^*$

$$\begin{aligned} H^\bullet \left(X, (\wedge^\bullet \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_X^{+1/2} \right) &= H^\bullet \left(X, (\wedge^\bullet \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right) \\ &= H^\bullet \left(X, (\wedge^\bullet \mathcal{E}^*) \otimes (\det \mathcal{E}^*)^{-1/2} \otimes K_X^{+1/2} \right) \end{aligned}$$

— manifest

- Should be implicit in elliptic genera

Leading term is proportional to

$$\begin{aligned} &\int \hat{A}(TX) \wedge \text{ch} \left((\det \mathcal{E})^{+1/2} \wedge_{-1}(\mathcal{E}^*) \right) \\ &= \int \text{td}(TX) \wedge \text{ch} \left(\wedge_{-1}(\mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right) \\ &= \sum_i (-)^i \chi \left((\wedge^i \mathcal{E}^*) \otimes (\det \mathcal{E})^{+1/2} \otimes K_X^{+1/2} \right) \end{aligned}$$

— matches

Sometimes we can perform a (pseudo-) topological twist.

These NLSM's have two anomalous global $U(1)$'s:

- a right-moving $U(1)_R$
- a canonical left-moving $U(1)$, rotating the phase of all left fermions, which becomes $U(1)_R$ on $(2,2)$ locus

If $\det \mathcal{E}^{\pm 1} \cong K_X$, then a nonanomalous $U(1)$ exists along which we can twist right & left moving fermions.

Possible twists....

A/2 model: Exists when $(\det \mathcal{E})^{-1} \cong K_X$

(on (2,2) locus, always possible; reduces to A model)

States: $H^\bullet(X, \wedge^\bullet \mathcal{E}^*)$

B/2 model: Exists when $\det \mathcal{E} \cong K_X$

(on (2,2) locus, requires $K_X^{\otimes 2} \cong \mathcal{O}_X$; reduces to B model)

States: $H^\bullet(X, \wedge^\bullet \mathcal{E})$

Exchanging $\mathcal{E} \leftrightarrow \mathcal{E}^*$ swaps the A/2, B/2 models.

(Physically, just a complex conjugation of left movers.)

Product structures

OPE rings in the $A/2$, $B/2$ models
= “quantum sheaf cohomology”

Physical relevance?

On the $(2,2)$ locus, in a perturbative heterotic compactification
on a CY 3-fold, say,

A model correlation f'ns & GW inv'ts encoded in $\overline{\mathbf{27}}^3$ couplings

B model correlation f'ns encoded in $\mathbf{27}^3$ couplings

Off the $(2,2)$ locus, Gromov-Witten inv'ts no longer relevant.

Mathematical GW computational tricks no longer apply.

No known analogue of periods, Picard-Fuchs equations.

New methods needed... and a few have been developed.

(A Adams, J Distler, R Donagi, J Guffin, S Katz, J McOrist, I Melnikov, R Plesser, ES,)

Review of quantum sheaf cohomology

Quantum sheaf cohomology is the heterotic version of quantum cohomology — defined by space + bundle.

Ex: ordinary quantum cohomology of \mathbb{P}^n

$$\mathbb{C}[x] / (x^{n+1} - q)$$

Compare: quantum sheaf cohomology of $\mathbb{P}^n \times \mathbb{P}^n$
with bundle

$$0 \rightarrow \mathcal{O} \oplus \mathcal{O} \xrightarrow{*} \mathcal{O}(1,0)^{n+1} \oplus \mathcal{O}(0,1)^{n+1} \rightarrow E \rightarrow 0$$

where

$$* = \begin{bmatrix} Ax & Bx \\ C\tilde{x} & D\tilde{x} \end{bmatrix} \quad x, \tilde{x} \text{ homog' coord's on } \mathbb{P}^n \text{'s}$$

is given by $\mathbb{C}[x,y] / (\det(Ax + By) - q_1, \det(Cx + Dy) - q_2)$

Check: When $E=T$, this becomes $\mathbb{C}[x,y] / (x^{n+1} - q_1, y^{n+1} - q_2)$

Review of quantum sheaf cohomology

Quantum sheaf cohomology

= OPE ring of the $A/2$ model

When does that OPE ring close into itself?

(2,2) susy **not** required.

For a SCFT, can use combination of

- worldsheet conformal invariance
- right-moving $N=2$ algebra

to argue closure on patches on moduli space.

(Adams-Distler-Ernebjerg, '05)

Review of quantum sheaf cohomology

Quantum sheaf cohomology

= OPE ring of the A/2 model

A model:

$$\text{Operators: } b_{i_1 \dots i_p \bar{i}_1 \dots \bar{i}_q} \chi^{\bar{i}_1} \cdots \chi^{\bar{i}_q} \cdots \chi^{i_1} \cdots \chi^{i_p} \leftrightarrow H^{p,q}(X)$$

A/2 model:

$$\text{Operators: } b_{\bar{i}_1 \dots \bar{i}_q a_1 \dots a_p} \psi_+^{\bar{i}_1} \cdots \psi_+^{\bar{i}_q} \lambda_-^{a_1} \cdots \lambda_-^{a_p} \leftrightarrow H^q(X, \wedge^p E^*)$$

On the (2,2) locus, A/2 reduces to A.

For operators, follows from

$$H^q(X, \wedge^p T^* X) = H^{p,q}(X)$$

Review of quantum sheaf cohomology

Quantum sheaf cohomology

= OPE ring of the A/2 model

Schematically:

A model: Classical contribution:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_X \omega_1 \wedge \cdots \wedge \omega_n = \int_X (\text{top-form})$$

$$\omega_i \in H^{p_i, q_i}(X)$$

A/2 model: Classical contribution:

$$\langle \mathcal{O}_1 \cdots \mathcal{O}_n \rangle = \int_X \omega_1 \wedge \cdots \wedge \omega_n$$

Now, $\omega_1 \wedge \cdots \wedge \omega_n \in H^{\text{top}}(X, \wedge^{\text{top}} E^*) = H^{\text{top}}(X, K_X)$

using the anomaly constraint $\det E^* \cong K_X$

Again, a top form, so get a number.

Review of quantum sheaf cohomology

To make this more clear, let's consider an

Example: classical sheaf cohomology on $\mathbb{P}^1 \times \mathbb{P}^1$

with gauge bundle E a deformation of the tangent bundle:

$$0 \rightarrow W^* \otimes \mathcal{O} \xrightarrow{*} \underbrace{\mathcal{O}(1,0)^2 \oplus \mathcal{O}(0,1)^2}_{Z^*} \rightarrow E \rightarrow 0$$

where $* = \begin{bmatrix} Ax & Bx \\ C\tilde{x} & D\tilde{x} \end{bmatrix}$ x, \tilde{x} homog' coord's on \mathbb{P}^1 's

and $W = \mathbb{C}^2$

Operators counted by $H^1(E^*) = H^0(W \otimes \mathcal{O}) = W$

n-pt correlation function is a map $\text{Sym}^n H^1(E^*) = \text{Sym}^n W \rightarrow H^n(\wedge^n E^*)$

OPE's = kernel

Plan: study map corresponding to classical corr' f'n

Review of quantum sheaf cohomology

Example: classical sheaf cohomology on $\mathbb{P}^1 \times \mathbb{P}^1$

with gauge bundle E a deformation of the tangent bundle:

$$0 \rightarrow W^* \otimes \mathcal{O} \xrightarrow{*} \underbrace{\mathcal{O}(1,0)^2 \oplus \mathcal{O}(0,1)^2}_{Z^*} \rightarrow E \rightarrow 0$$

where $* = \begin{bmatrix} Ax & Bx \\ C\tilde{x} & D\tilde{x} \end{bmatrix}$ x, \tilde{x} homog' coord's on \mathbb{P}^1 's

and $W = \mathbb{C}^2$

Since this is a rk 2 bundle, classical sheaf cohomology defined by products of 2 elements of $H^1(E^*) = H^0(W \otimes \mathcal{O}) = W$.

So, we want to study map $H^0(\text{Sym}^2 W \otimes \mathcal{O}) \rightarrow H^2(\wedge^2 E^*) = \text{corr}' \text{ f'n}$

This map is encoded in the resolution

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

Review of quantum sheaf cohomology

Example: classical sheaf cohomology on $\mathbb{P}^1 \times \mathbb{P}^1$

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

Break into short exact sequences:

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow S_1 \rightarrow 0$$

$$0 \rightarrow S_1 \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

Examine second sequence:

$$\text{induces } \begin{array}{ccccccc} H^0(Z \otimes W) & \rightarrow & H^0(\text{Sym}^2 W \otimes \mathcal{O}) & \xrightarrow{\delta} & H^1(S_1) & \rightarrow & H^1(Z \otimes W) \\ \searrow & & & & & & \searrow \\ & & 0 & & & & 0 \end{array}$$

Since Z is a sum of $\mathcal{O}(-1,0)$'s, $\mathcal{O}(0,-1)$'s,

$$\text{hence } \delta : H^0(\text{Sym}^2 W \otimes \mathcal{O}) \xrightarrow{\sim} H^1(S_1) \quad \text{is an iso.}$$

Next, consider the other short exact sequence at top....

Review of quantum sheaf cohomology

Example: classical sheaf cohomology on $\mathbb{P}^1 \times \mathbb{P}^1$

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

Break into short exact sequences:

$$0 \rightarrow S_1 \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

$$\delta : H^0(\text{Sym}^2 W \otimes \mathcal{O}) \xrightarrow{\sim} H^1(S_1)$$

Examine other sequence:

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow S_1 \rightarrow 0$$

$$\text{induces } H^1(\wedge^2 Z) \rightarrow H^1(S_1) \xrightarrow{\delta} H^2(\wedge^2 E^*) \rightarrow H^2(\wedge^2 Z) \rightarrow 0$$

Since Z is a sum of $\mathcal{O}(-1,0)$'s, $\mathcal{O}(0,-1)$'s,

$$H^2(\wedge^2 Z) = 0 \quad \text{but} \quad H^1(\wedge^2 Z) = \mathbb{C} \oplus \mathbb{C}$$

and so $\delta : H^1(S_1) \rightarrow H^2(\wedge^2 E^*)$ has a 2d kernel.

Now, assemble the coboundary maps....

Review of quantum sheaf cohomology

Example: classical sheaf cohomology on $\mathbb{P}^1 \times \mathbb{P}^1$

$$0 \rightarrow \wedge^2 E^* \rightarrow \wedge^2 Z \rightarrow Z \otimes W \rightarrow \text{Sym}^2 W \otimes \mathcal{O} \rightarrow 0$$

Now, assemble the coboundary maps.....

A classical (2-pt) correlation function is computed as

$$H^0(\text{Sym}^2 W \otimes \mathcal{O}) \xrightarrow{\tilde{\delta}} H^1(S_1) \xrightarrow{\delta} H^2(\wedge^2 E^*)$$

where the right map has a 2d kernel, which one can show is generated by

$$\det(A\psi + B\tilde{\psi}), \quad \det(C\psi + D\tilde{\psi})$$

where A, B, C, D are four matrices defining the def' E , and $\psi, \tilde{\psi}$ correspond to elements of a basis for W .

Classical sheaf cohomology ring:

$$\mathbb{C}[\psi, \tilde{\psi}] / (\det(A\psi + B\tilde{\psi}), \det(C\psi + D\tilde{\psi}))$$

Review of quantum sheaf cohomology

Quantum sheaf cohomology

= OPE ring of the $A/2$ model

Instanton sectors have the same form,
except X replaced by moduli space M of instantons,
 E replaced by induced sheaf F over moduli space M .

Must compactify M ,
and extend F over compactification divisor.

$$\left. \begin{array}{l} \wedge^{\text{top}} E^* \cong K_X \\ \text{ch}_2(E) = \text{ch}_2(TX) \end{array} \right\} \xRightarrow{\text{GRR}} \wedge^{\text{top}} F^* \cong K_M$$

Within any one sector, can follow the same method just outlined....

Review of quantum sheaf cohomology

In the case of our example,
one can show that in a sector of instanton degree (a,b) ,
the 'classical' ring in that sector is of the form

$$\text{Sym}^{\bullet} \mathbf{W} / (Q^{a+1}, \tilde{Q}^{b+1})$$

where $Q = \det(A\psi + B\tilde{\psi}), \quad \tilde{Q} = \det(C\psi + D\tilde{\psi})$

Now, OPE's can relate correlation functions in different instanton degrees, and so, should map ideals to ideals.

To be compatible with those ideals,

$$\langle \mathcal{O} \rangle_{a,b} = q^{a'-a} \tilde{q}^{b'-b} \langle \mathcal{O} Q^{a'-a} \tilde{Q}^{b'-b} \rangle_{a',b'}$$

for some constants $q, \tilde{q} \Rightarrow$ OPE's $Q = q, \tilde{Q} = \tilde{q}$

— quantum sheaf cohomology rel'ns

Review of quantum sheaf cohomology

General result:

(Donagi, Guffin, Katz, ES, '11)

For any toric variety, and any def' E of its tangent bundle,

$$0 \rightarrow W^* \otimes \mathcal{O} \rightarrow \underbrace{\bigoplus \mathcal{O}(\vec{q}_i)}_{Z^*} \rightarrow E \rightarrow 0$$

the chiral ring is

$$\prod_{\alpha} (\det M_{(\alpha)})^{Q_{\alpha}^a} = q_a$$

where the M 's are matrices of chiral operators built from $*$.

Review of quantum sheaf cohomology

So far, I've outlined mathematical computations of quantum sheaf cohomology, but GLSM-based methods also exist:

- Quantum cohomology ((2,2)): Morrison-Plesser '94
- Quantum sheaf cohomology ((0,2)): McOrist-Melnikov '07, '08

Briefly, for (0,2) case:

One computes quantum corrections to effective action of form

$$L_{\text{eff}} = \int d\theta^+ \sum_a Y_a \log \left[\prod_{\alpha} (\det M_{(\alpha)})^{Q_{\alpha}^a} / q_a \right]$$

from which one derives $\prod_{\alpha} (\det M_{(\alpha)})^{Q_{\alpha}^a} = q_a$

— these are q.s.c. rel'ns

— match math' computations

Review of quantum sheaf cohomology

State of the art: computations on toric varieties

To do: compact CY's

Intermediate step: Grassmannians (work in progress)

Briefly, what we need are better computational methods.

Conventional GW tricks seem to revolve around idea that A model is independent of complex structure, not necessarily true for $A/2$.

- [McOrist-Melnikov '08](#) have argued an analogue for $A/2$
- Despite attempts to check ([Garavuso-ES '13](#)), still not well-understood

So far, I've (secretly) been talking about abelian GLSM's.

Next, let's turn to nonabelian GLSMs:

- Dualities in 2d and their geometry
- Tests of Gaiotto-Gukov-Putrov triality

In 2d theories, dualities often have a purely geometric understanding.

Trivial example:

$U(k)$ gauge theory,
 n chiral multiplets



NLSM on $G(k,n)$

=

$U(n-k)$ gauge theory,
 n chiral multiplets



NLSM on $G(n-k,n)$

Another example, in 2d, (2,2) susy:

U(k) gauge group,

matter: n chirals in fund' \mathbf{k} , $n > k$,

A chirals in antifund' \mathbf{k}^* , $A < n$

Seiberg
 \longleftrightarrow
 Benini-Cremonesi '12

U(n-k) gauge group,

matter: n chirals Φ in fund' \mathbf{k} ,

A chirals P in antifund' \mathbf{k}^* ,

nA neutral chirals M,

superpotential: $W = M \Phi P$



NLSM on $\text{Tot}(S^A \rightarrow G(k, n))$
 $= (\mathbb{C}^{kn} \times \mathbb{C}^{kA}) // GL(k)$



...



$= \text{Tot}((Q^*)^A \rightarrow G(n-k, n))$




...

Build physics for RHS using $0 \rightarrow S \xrightarrow{\Phi} \mathcal{O}^n \rightarrow Q \rightarrow 0$
 & discover the upper RHS.

So, 2d analogue of Seiberg duality has geometric description.

Another example, in 2d, (2,2) susy:



The physical duality implied at top relates abelian & nonabelian gauge theories, which in 4d for ex would be surprising.

Further examples

((2,2) susy)

U(2) gauge theory, \longleftrightarrow U(n-2)xU(1) gauge theory,
 n chirals in fundamental \longleftrightarrow n chirals X in fundamental of U(n-2),
 n chirals P in antifundamental of U(n-2)

(n choose 2) chirals $z_{ij} = -z_{ji}$
 each of charge +1 under U(1),

$$W = \text{tr PAX}$$



$G(2,n) = \text{rank 2 locus of } n \times n \text{ matrix } A \text{ over } \mathbb{P}^{\binom{n}{2}-1}$

$$A(z_{ij}) = \begin{bmatrix} z_{11} = 0 & z_{12} & z_{13} & \dots \\ z_{21} = -z_{12} & z_{22} = 0 & z_{23} & \dots \\ z_{31} = -z_{13} & z_{32} = -z_{23} & z_{33} = 0 & \dots \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

(using description of Pfaffians of
 Hori '11, Jockers et al '12)

In this fashion, straightforward to generate examples;
 let's move on.....

Triality

((0,2) susy)

(Gadde-Gukov-Putrov '13-'14)

GGP proposed that *triples* of (0,2) GLSM's might flow to the same IR fixed point.

In terms of lower-energy NLSM's, the theories are

Gauge bundle \longrightarrow Target space

$$S^A \oplus (Q^*)^{2k+A-n} \oplus (\det S^*)^2 \longrightarrow G(k, n)$$

$$S^{2k+A-n} \oplus (Q^*)^n \oplus (\det S^*)^2 \longrightarrow G(n-k, A)$$

$$S^n \oplus (Q^*)^A \oplus (\det S^*)^2 \longrightarrow G(A-n+k, 2k+A-n)$$

related by permuting 3 of flavor symmetries.

Susy unbroken iff geometric description above valid.

Gadde-Gukov-Putrov triality ('13) ((0,2) susy)

bundle

space

$$S^A \oplus (Q^*)^{2k+A-n} \rightarrow G(k, n) \quad \dots \text{phase} \dots \quad (S^*)^A \oplus (Q^*)^n \rightarrow G(k, 2k + A - n)$$

$$\updownarrow =$$

$$(Q^*)^A \oplus S^{2k+A-n} \rightarrow G(n-k, n) \quad \dots \text{phase} \dots \quad (Q^*)^n \oplus (S^*)^{2k+A-n} \rightarrow G(n-k, A)$$

$$\updownarrow =$$

$$S^n \oplus (Q^*)^A \rightarrow G(A-n+k, 2k+A-n) \quad \dots \text{phase} \dots \quad (S^*)^n \oplus (Q^*)^{2k+A-n} \rightarrow G(A-n+k, A)$$

$$\updownarrow =$$


$$(Q^*)^n \oplus S^A \rightarrow G(k, 2k + A - n) \quad \dots \text{phase} \dots \quad (Q)^{2k+A-n} \oplus S^A \rightarrow G(k, n)$$

For brevity, I've omitted writing out the (0,2) gauge theory.

Utilizes another duality: $\text{NLSM}(X, E) = \text{NLSM}(X, E^*)$

Triality predicts

(0,2) NLSM's:



IR fixed point:

IR SCFT = (left-moving Kac-Moody) \otimes (right-moving Kazama-Suzuki)

UV global $SU(n) \times SU(A) \times SU(2k + A - n) \times SU(2)$

(present in GLSM & each NLSM)


enhanced in IR to affine

$SU(n)_{k+A-n} \times SU(A)_k \times SU(2k + A - n)_{n-k} \times SU(2)_1$

Chiral states should live in integrable reps of affine algebras.

Let's check triality, using chiral rings.

(0,2) NLSM's:



IR fixed point:

Plan: Compute chiral states in each theory and compare.

Alas, not quite so simple.....

Subtleties in comparing chiral states:

- Q^* -cohomology in large-radius $(0,2)$ NLSM invariant under perturbative corrections, but, here RG flow goes to strong coupling — states might enter/leave.

We'll see exactly that —

not all states will match between different presentations, but, states that don't match, shouldn't be in IR either.

- Chiral ring computations in 2d KS models not under good control; Lie algebra cohomology is part of answer.

We'll focus on comparing states across UV presentations, then, merely outline in general terms how form of Lie algebra cohomology is appropriate.

Example 1:

$r \gg 0$:

$$\mathcal{E} = U \otimes S + V \otimes Q^* + W \otimes \det S^* \longrightarrow \mathbb{P}\tilde{V}^* = \mathbb{P}^2$$

$r \ll 0$:

$$\mathcal{E} = U \otimes S^* + \tilde{V} \otimes Q^* + W \otimes \det S \longrightarrow \mathbb{P}V^* = \mathbb{P}^1$$

$$U = \mathbb{C}^3, \quad V = \mathbb{C}^2, \quad W = \mathbb{C}^2, \quad \tilde{V} = \mathbb{C}^3$$

Let's compare states in these two phases
 (= 2 of 3 triality-related geometries)....

Example 1:

$r \gg 0$:

$$\mathcal{E} = U \otimes S + V \otimes Q^* + W \otimes \det S^* \longrightarrow \mathbb{P}\tilde{V}^* = \mathbb{P}^2$$

Compute states:

$$H^\bullet(\mathbb{P}^2, (\wedge^\bullet \mathcal{E}) \otimes (\det \mathcal{E})^{-1/2} \otimes K_{\mathbb{P}^2}^{+1/2})$$

Global symmetries:

$SU(U) \times SU(V) \times SU(W)$ manifest — acts on bundle

$SU(\tilde{V})$ also present:

Compute sheaf cohomology with Bott-Borel-Weil, which gives sheaf cohomology as reps of $U(\tilde{V})$.

These computations are an application of Bott-Borel-Weil,
so, brief overview:

For a bundle \mathcal{E}_ξ on G/P defined by rep' ξ of P ,
 $H^\bullet(G/P, \mathcal{E}_\xi)$ is naturally a rep' of G .

For Grassmannians,

compute $H^\bullet(G(k, n), K_{(a_1, \dots, a_k)} S^* \otimes K_{(b_1, \dots, b_{n-k})} Q^*)$:

(a_1, \dots, a_k) rep' of $U(k)$ $a_1 \geq a_2 \geq \dots \geq a_k$

(b_1, \dots, b_{n-k}) rep' of $U(n-k)$ $b_1 \geq b_2 \geq \dots \geq b_{n-k}$

'Mutate' $(a_1, \dots, a_k, b_1, \dots, b_{n-k})$ to (c_1, \dots, c_n) rep of $U(n)$

$H^\bullet(G(k, n), K_{(a_1, \dots, a_k)} S^* \otimes K_{(b_1, \dots, b_{n-k})} Q^*) = K_{(c_1, \dots, c_n)} V^*$

for \bullet = number of mutations, & zero in other degrees.

Bott-Borel-Weil, cont'd

$$\text{Ex: } H^\bullet(G(k, \tilde{V}^*), U \otimes S^*)$$

$$= U \otimes H^\bullet(G(k, \tilde{V}^*), K_{(1,0,\dots,0)} S^* \otimes K_{(0,0,\dots,0)} Q^*)$$

$$= U \otimes K_{(1,0,\dots,0)} \tilde{V} \delta^{\bullet,0} = U \otimes \tilde{V} \delta^{\bullet,0}$$

Constraints on results:

- Invariance under Serre duality

$$H^\bullet(X, \mathcal{E}) = H^{\dim - \bullet}(X, \mathcal{E}^* \otimes K_X)^*$$

Should map state spectrum into itself,
dualizing representation.

- Integrability of representations

GGP triality predicts that states should live
in 'integrable' rep's.

$SU(n)_k$: integrable reps have Young tableaux of width $\leq k$

Let's look at some states in the first example....

Examples of states shared between two phases:



| $SU(3) \times SU(2) \times SU(2) \times SU(3)$ | $U(1)$ |
|--|------------------|
| $(1,1,1,1)$ | $(+3,0,-3)$ |
| $(1,1,2,3)$ | $(+2,-1/2,-3/2)$ |
| $(1,1,1,3^*)$ | $(+1,+2,-3)$ |
| $(3,2,1,1)$ | $(+2,0,-2)$ |
| $(3,1,2,1)$ | $(-2,-3/2,-1/2)$ |
| $(1,2,2,3^*)$ | $(+1,+1/2,-3/2)$ |
| $(3,1,1,3)$ | $(+1,+1,-2)$ |
| ... | ... |
| $(3^*,1,1,3^*)$ | $(-1,-1,+2)$ |
| $(1,2,2,3)$ | $(-1,-1/2,+3/2)$ |
| $(3^*,1,2,1)$ | $(-2,+3/2,+1/2)$ |
| $(3^*,2,1,1)$ | $(-2,0,+2)$ |
| $(1,1,1,3)$ | $(-1,-2,+3)$ |
| $(1,1,2,3^*)$ | $(-2,+1/2,+3/2)$ |
| $(1,1,1,1)$ | $(-3,0,+3)$ |


Serre
duals




Integrable reps of $SU(3)_1 \times SU(2)_2 \times SU(2)_1 \times SU(3)_1$

Non-shared states in $r \gg 0$ phase:

| wedge | coh' | degree | SU(3)xSU(2)xSU(2)xSU(3) | U(1) |
|-------|------|--------|-------------------------|-----------|
| 2 | 0 | | (1,1,1,6) | (+1,-1,0) |
| 3 | 0 | | (1,2,1,8) | (0,0,0) |
| 4 | 0 | | (1,1,1,6*) | (-1,+1,0) |
| 5 | 2 | | (1,1,1,6) | (+1,-1,0) |
| 6 | 2 | | (1,2,1,8) | (0,0,0) |
| 7 | 2 | | (1,1,1,6*) | (-1,+1,0) |


 Non-integrable rep' of $SU(3)_1 \times SU(2)_2 \times SU(2)_1 \times SU(3)_1$


 Serre duals
 Matching rep'

- All states come in Serre dual pairs
- Rep's are non-integrable — should not survive to IR
 - States come in pairs with matching rep's (so cancel out of elliptic genera)

Example 2:

$r \gg 0$:

$$\mathcal{E} = U \otimes S + V \otimes Q^* + W \otimes \det S^* \longrightarrow \mathbb{P}\tilde{V}^* = \mathbb{P}^1$$

$r \ll 0$:

$$\mathcal{E} = U \otimes S^* + \tilde{V} \otimes Q^* + W \otimes \det S \longrightarrow \mathbb{P}V^* = \mathbb{P}^3$$

$$U = \mathbb{C}^4, \quad V = \mathbb{C}^4, \quad W = \mathbb{C}^2, \quad \tilde{V} = \mathbb{C}^2$$

Let's compare states in these two phases
 (= 2 of 3 triality-related geometries)....

Examples of states shared between two phases:




| SU(4)xSU(4)xSU(2)xSU(2) | U(1) |
|--------------------------------|-------------|
| (1,1,1,3) | (+4,0,-4) |
| (4,1,1,2) | (+4,-1,-3) |
| (1,4,1,2) | (-3,+1,-4) |
| (1,1,2,4) | (+2,0,-2) |
| (6,1,1,1) | (+4,-2,-2) |
| (4,4,1,1) | (+3,0,-3) |
| (4,1,2,3) | (+2,-1,-1) |
| ... | ... |
| (4*,1,2,3) | (-2,+1,+1) |
| (4*,4*,1,1) | (-3,0,+3) |
| (6,1,1,1) | (-4,+2,+2) |
| (1,1,2,4) | (-2,0,+2) |
| (1,4*,1,2) | (-3,-1,+4) |
| (4*,1,1,2) | (-4,+1,+3) |
| (1,1,1,3) | (-4,0,+4) |


Serre
duals

Integrable reps of $SU(4)_1 \times SU(4)_1 \times SU(2)_1 \times SU(2)_3$

Non-shared states in $r \ll 0$ phase:

| wedge | coh' | degree | SU(4)xSU(4)xSU(2)xSU(2) | U(1) |
|-------|------|--------|-------------------------|-----------|
| 4 | 0 | | (1,10,1,1) | (+2,-2,0) |
| 4 | 3 | | (1,10,1,1) | (+2,-2,0) |
| 5 | 0 | | (1,20,1,2) | (+1,-1,0) |
| ... | ... | | ... | ... |
| 7 | 3 | | (1,20,1,2) | (-1,+1,0) |
| 8 | 0 | | (1,10*,1,1) | (-2,+2,0) |
| 8 | 3 | | (1,10*,1,1) | (-2,+2,0) |


 Non-integrable rep' of $SU(4)_1 \times SU(4)_1 \times SU(2)_1 \times SU(2)_3$



- All states come in Serre dual pairs
- Rep's are non-integrable — should not survive to IR
 - States come in pairs with matching rep's (so cancel out of elliptic genera)

We've seen that the between geometries that should flow to same fixed point, the chiral states don't all match,
but,

the ones that don't, also have nonintegrable reps,
and make no net contribution to refined elliptic genera.

We believe that they get a mass and disappear from RG flow.

The fact that the remaining states are both

- shared between phases, and
- in integrable reps of proposed IR symmetry algebras,
serves as a nontrivial check of triality.

How in principle might these UV sheaf cohomology groups relate, in general, to the IR states?

In IR, expect states \sim Lie algebra cohomology.

[*roughly* — correspondence incomplete]

(W Lerche,
private communication)

How is that related?

We won't pursue this in detail, but, want to observe that another flavor of BBW provides the missing link:

sheaf
cohomology

$$H^\bullet(G/P, \mathcal{E}_\xi)_\lambda = H^\bullet(\mathfrak{n}, V_\lambda)_\xi$$


Lie algebra
cohomology

λ a representation of G

ξ a representation of P

$$\mathfrak{p} = (\text{Levi}) + \mathfrak{n}$$

Summary:

- Chiral states in 2d (0,2) NLSM's
- Product structures in chiral rings in $A/2$, $B/2$ twists:
quantum sheaf cohomology
- Nonabelian GLSMs:
 - Dualities in 2d and their geometry
 - Tests of Gadde-Gukov-Putrov triality 

Thank you for your time!